ScienceDirect

Brain Research

Volume 1624, 22 October 2015, Pages 232-238

Research Report

Deleterious impacts of a 900-MHz electromagnetic field on hippocampal pyramidal neurons of 8-week-old Sprague Dawley male rats

Arzu Şahin [a], Ali Aslan [a], Orhan Baş [b], Ayşe İkinci [c], Cansu Özyılmaz [c], Osman Fikret Sönmez [d], Serdar Çolakoğlu [e], Ersan Odacı [c]

Show more

Share Cite

https://doi.org/10.1016/j.brainres.2015.07.042

Get rights and content

Highlights

- Rat hippocampus was investigated following exposure to 900MHz electromagnetic field.
- The number of pyramidal neurons in the cornu ammonis of the hippocampus was estimated.
- Cells with abnormal and black cytoplasm among the normal neurons were observed.
- Largest lateral ventricles were observed in the electromagnetic field group sections.

- Pyramidal neuron loss in the cornu ammonis of electromagnetic field group was showed.

Abstract

Children are at potential risk due to their intense use of mobile phones. We examined 8-week-old rats because this age of the rats is comparable with the preadolescent period in humans. The number of pyramidal neurons in the cornu ammonis of the Sprague Dawley male rat (8-weeks old, weighing 180–250 g) hippocampus following exposure to a 900 MHz (MHz) electromagnetic field (EMF) were examined. The study consisted of control (CN-G), sham exposed (SHM-EG) and EMF exposed (EMF-EG) groups with 6 rats in each. The EMF-EG rats were exposed to 900 MHz EMF (1 h/day for 30 days) in an EMF jar. The SHM-EG rats were placed in the EMF jar but not exposed to the EMF (1 h/day for 30 days). The CN-G rats were not placed into the exposure jar and were not exposed to the EMF during the study period. All animals were sacrificed at the end of the experiment, and their brains were removed for histopathological and stereological analysis. The number of pyramidal neurons in the cornu ammonis of the hippocampus was estimated on Cresyl violet stained sections of the brain using the optical dissector counting technique. Histopathological evaluations were also performed on these sections. Histopathological observation showed abundant cells with abnormal, black or dark blue cytoplasm and shrunken morphology among the normal pyramidal neurons. The largest lateral ventricles were observed in the EMF-EG sections compared to those from the other groups. Stereological analyses showed that the total number of pyramidal neurons in the cornu ammonis of the EMF-EG rats was significantly lower than those in the CN-G ($p<0.05$) and the SHM-EG ($p<0.05$). In conclusion, our results suggest that pyramidal neuron loss and histopathological changes in the cornu ammonis of 8-week-old male rats may be due to the 900-MHz EMF exposure.

Introduction

Several studies have reported that the electromagnetic field (EMF) emitted from natural or non-natural sources has severe effects on human health and that these can give rise to significant pathologies in various tissues and organs (Davis et al., 2013, Gultekin and Moeller, 2013). The most popular communication device today is the mobile phone, an important source of EMF. The effects of EMF on human health established by mobile phones is therefore the subject of significant research and debate among scientists (Hancı et al., 2013, Hancı et al., 2015, İkinci et al., 2013, Türedi et al., 2014, Topal et al., 2015). Mobile

phones are the source of EMF used in the greatest proximity to the human body, and the field established by them includes organs and tissues with vital functions such as the brain, cerebellum, eye and hearing organs. Some scientists maintain that these effects can have serious pathological outcomes affecting human health (Davis et al., 2013), while others maintain the opposite (Inskip et al., 2010).

Gultekin and Moeller (2013) recently showed that currently used cell phones can produce hotspots in living brain tissue or brain-related tissue/organs. In another study, Volkow et al. (2011) reported that 50 min of mobile phone use produces a significant change in glucose metabolism in the area of the human brain that absorbs most of the radiation. Although the cause of the marked increase in pathological conditions in humans such as brain tumors or skin cancer incidence is unknown (Corle et al., 2012, Inskip et al., 2010, Poulsen et al., 2013), there is nevertheless concern that cell phones can trigger biological effects and that several decades of cell phone use in an individual is a probable carcinogen (Davis et al., 2013).

Children's brains are more affected by EMF than the adult brains when speaking on mobile phones. This is because the brain membrane, skull bones and components associated with those bones that protect central nervous system (CNS) structures inside the skull against external agents have not yet achieved the thickness and morphological characteristics of those in adulthood. In the light of studies reporting that exposure, even in adulthood, can cause significant pathological changes in CNS structures (Bas et al., 2009a, Bas et al., 2009b, Odaci et al., 2008, Sonmez et al., 2010), exposure to EMF in childhood may well affect the child brain and related structures and the development thereof.

The CNS of rodents (i.e. rats) tested must be comparable to that in the same stage of development in humans, regardless of whether it is tested during the fetal, prenatal or postnatal periods (Dobbing, 1970, Dobbing and Sands, 1973, Jacobson, 1991, Rodier, 1980). We examined the brains of 8-week-old rats after exposure to 900-MHz megahertz (MHz) EMF because that age is comparable with the human preadolescent period (Baş et al., 2009b; Odaci et al., 2004, Odaci et al., 2008). In addition to histopathological examination, we also used the stereological optical dissector method for neuron number estimation because it represents an unbiased means of estimating the total neuron numbers.

Access through your organization

Check access to the full text by signing in through your organization.

Access through **Haverford College**

Section snippets

Physical examination and weights

Physical examination revealed no skeletal anomaly or pathological findings in the CN-G, SHM-EG or EMF-EG rats. No significant difference was determined among the groups in terms of rats′ body or brain weights at the end of the experiment ($p>0.05$), (Table 1). …

Pyramidal neuron numbers

The total number of pyramidal neurons in the cornu ammonis of the hippocampus was significantly lower in the EMF-EG rats compared to the CN-G and the SHM-EG ($p<0.05$). There was no significant difference in total pyramidal neuron numbers …

Discussion

The number of mobile phone subscriptions worldwide is approaching the number of people on Earth, reaching almost 7 billion by the end of 2014 (ITU, 2014). The mobile phone has become the favored communication tool for the majority of children and teenagers (Lenhart et al., 2010), and these are at potential risk due to their intense use of mobile phones. Their brain and its associated structures are therefore affected by EMF while speaking on the mobile phones. We therefore elected 8-week-old …

Animals and groups

Eighteen 8-week-old healthy Sprague Dawley male rats, with initial weights of 180–250g, were obtained from the Experimental Animals Surgical Research and Application Center of Karadeniz Technical University (EXASURAC-KTU). Experiments and procedures were performed in accordance with the Guide for the Care and Use of Laboratory Animals published by the U.S. National Institutes of Health. Ethical approval was obtained from the Karadeniz Technical University Animal Care and Ethics Committee. …

Recommended articles

References (39)

O. Bas *et al.*
900MHz electromagnetic field exposure affects qualitative and quantitative features of hippocampal pyramidal cells in the adult female rat
Brain Res. (2009)

D.L. Davis *et al.*
Swedish review strengthens grounds for concluding that radiation from cellular and cordless phones is a probable human carcinogen
Pathophysiology (2013)

H. Hancı *et al.*
The effect of prenatal exposure to 900-MHz electromagnetic field on the 21-old-day rat testicle
Reprod. Toxicol. (2013)

J. Hausdorf *et al.*
Extracorporeal shockwave application to the distal femur of rabbits diminishes the number of neurons immunoreactive for substance P in dorsal root ganglia L5
Brain Res. (2008)

E. Odaci *et al.*
Effects of prenatal exposure to a 900MHz electromagnetic field on the dentate gyrus of rats: a stereological and histopathological study
Brain Res. (2008)

E. Odaci *et al.*
Prenatal diclofenac sodium administration increases the number of Purkinje cells in female rats: a stereological study
Int. J. Dev. Neurosci. (2010)

O.F. Sonmez *et al.*
Purkinje cell number decreases in the adult female rat cerebellum following exposure to 900MHz electromagnetic field
Brain Res. (2010)

E. Tirelli *et al.*
Ontogenesis of behavioral sensitization and conditioned place preference induced by psychostimulants in laboratory rodents
Neurosci. Biobehav. Rev. (2003)

A.T. Tunc *et al.*

Inhibitory effect of pinealectomy on the development of cerebellar granule cells in the chick: a stereological study

Brain Res. (2007)

O. Bas *et al.*

Chronic prenatal exposure to the 900MHz electromagnetic field induces pyramidal cell loss in the hippocampus of newborn rats

Toxicol. Ind. Health (2009)

View more references

Cited by (28)

Effect of 2400 MHz mobile phone radiation exposure on the behavior and hippocampus morphology in Swiss mouse model: Effect of 4G mobile phone radiation on the brain

2022, Saudi Journal of Biological Sciences

Citation Excerpt :

…Most noticeable behavioral and cognitive disorders of the central nervous system identified in both people and animals regarding long-term smartphone radiation exposure (Babadi-Akashe et al., 2014; Söderqvist et al., 2015; Zhao et al., 2012). Exposure to EMF has indeed been documented to have a variety of neurological consequences, such as decreased number of Cornu Ammonis (CA) pyramidal and Purkinje cells (Aslan et al., 2017; Baş et al., 2013;Odaci et al., 2008; Şahin et al., 2015), increased dark cell cytoplasm (Salford et al., 2003), behavioral changes (Ikinci et al., 2013) and concentration disturbances (Durusoy et al., 2017). Many research focused on behavioral problems caused by RF-EMR exposure, including learning and memory loss in rats (Lai et al., 1994; Wang and Lai, 2000).…

Show abstract

Absorption of 5G Radiation in Brain Tissue as a Function of Frequency, Power and Time ↗

2020, IEEE Access

Hippocampal lipidome and transcriptome profile alterations triggered by acute exposure of mice to GSM 1800 MHz mobile phone radiation: An exploratory study

↗

2018, Brain and Behavior

Oxidative stress response in SH-SY5Y cells exposed to short-term 1800MHz radiofrequency radiation ↗

2018, Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering

EUROPAEM EMF Guideline 2016 for the prevention, diagnosis and treatment of EMF-related health problems and illnesses ↗

2016, Reviews on Environmental Health

Electromagnetic radiation of mobile telecommunication antennas affects the abundance and composition of wild pollinators ↗

2016, Journal of Insect Conservation

View all citing articles on Scopus ↗

View full text

Copyright © 2015 Elsevier B.V. All rights reserved.

All content on this site: Copyright © 2025 or its licensors and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the relevant licensing terms apply.